

Assessment of educated people awareness level and sources about COVID-19

Marwa O. Elgendy^{1, 2}, Haitham Saeed³, Heba A. Abou-Taleb⁴

Correspondence:

Marwa O. Elgendy
Department of Clinical Pharmacy,
Beni-Suef University Hospitals,
Faculty of Medicine, Beni-Suef
University, Beni-Suef, Egypt
Department of Clinical Pharmacy,
Faculty of Pharmacy, Nahda
University (NUB), Egypt.
marwaosamaelgendy@yahoo.com

1 Department of Clinical Pharmacy,
Beni-Suef University Hospitals,
Faculty of Medicine, Beni-Suef
University, Beni-Suef, Egypt
2 Department of Clinical Pharmacy,
Faculty of Pharmacy, Nahda
University (NUB), Egypt.
3 Clinical Pharmacy Department,
Faculty of Pharmacy, Beni-Suef
University, Beni-Suef 62521, Egypt
4 Department of Pharmaceutics
and Industrial Pharmacy, Faculty of
Pharmacy, Merit University (MUE),
Sohag 82755, Egypt

Volume number 1
Issue number 1
Pages 19-27

10.61466/ijcmr1010004

Received: 11.07.2023
Accepted: 18.08.2023
Published: 20.08.2023
Online: 20.08.2023

Abstract

Background:

Measurement of educated people knowledge about COVID-19 is important to enhance health measures to reduce the rate of infection. Also, knowing the most usable source of data for the public is important point for enhancing the public awareness about the disease. Hence, this study aimed to evaluate the awareness and the practices related to COVID-19 among educated people.

Methods:

360 participant received a questionnaire that was divided into several parts aimed at collecting data about participant awareness regarding COVID-19. Data collected using a designed questionnaire; to evaluate the knowledge related to COVID-19. The questionnaire was classified into 6 parts covering all aspects of COVID-19 awareness data. The first part was to evaluate the respondent's general knowledge [6 items]; the second was to evaluate symptoms knowledge [1 item]; the third was to evaluate transmission knowledge [2 items]; the fourth was to evaluate preventive knowledge [4 items]; the fifth was to evaluate treatment knowledge [2 items] and the sixth was indicated the main source of gathering information about the disease [1 item].

Results:

A total of 360 participants participated, 91% of them know the main clinical symptoms of COVID-19. 86% believed that following preventive measures (the etiquette of sneezing, coughing, personal hygiene or wearing a mask) reduce infection transmission. More than 90 % of participants used the social media for gathering information about the disease, while highly educated (Master and PhD holders) used the published articles as a source for their knowledge, however they represent a few percent of the participant.

Conclusion:

Overall, the study participants had good knowledge about COVID-19, however, more efforts are needed to maximize the awareness of the public. Further clinical study is required to evaluate the effect of such good knowledge on decreasing the infection rate.

Keywords: COVID-19; highly educated public awareness; Egypt

Introduction

Respiratory viral diseases which started and spread during the current century were a result of coronaviruses infections. ^{1, 2} one of the theses attacks affected all over the world and lead to more than million death, COVID-19 is viral respiratory infection, which is pandemic all over the world ^{3, 4}. The spread of COVID-19 infections depend on the transmission of droplets between people, hence the direct contact with confirmed cases is the big source for catching the infection ⁵⁻⁷. The diseases does not affect all ages with the same strength, elderly are more affected and high percent of them become hospitalized and need mechanical ventilation ^{8, 9}. At the start of the pandemic, the symptoms of the disease were mainly related to the respiratory system such as cough, fever and breathing difficulties, which could progress to result in acute respiratory failure syndrome or multi organ failure, and death ^{6, 10}. Now more non-respiratory symptoms were recorded such as gastro-intestinal related symptoms (diarrhea and vomiting) which were confirmed by presence of the viral genetic material in stool samples. ^{11, 12} Loss of ability to smell or taste your food could be

related to COVID-19 infection.^{13, 14}

According to the nature of COVID-19 spread, improving the public awareness about the importance of social distancing and self-protective tools is the essential element in limiting the spread of the infectious diseases^{15, 16}. Proper education to the public could be helpful in reducing the rate of infection also in early diagnosis and treatment of the disease^{17, 18}. Limiting the spread of the infection has a lot of benefits not only by decreasing the number of cases or death but also it has financial aspects. The higher the spread rate, the lower the possibility of good medical care because of the possible medical shortage that will result of receiving large number of subjects that exceeded the hospital capacities.^{19, 20}

To reduce the spreading of the infection, governments' Ministry of Health develops awareness and educational messages to educate the public about this virus.²¹

This study aimed to evaluate the awareness and the practices related to COVID-19 among highly educated people and the most suitable way for increasing the public awareness.

Methods

All of the experiments described here were conducted with permission from the Institutional Ethics Committee at Beni-Suef University Hospitals. A study was conducted for the period on highly educated people to assess their knowledge and awareness about COVID-19. All subjects participated in the study received a questionnaire which were divided into seven sections. The first section was oriented to collect demographic data from subjects (gender, age in year, marital status, current place of residence (city or country), and educational level). The second section was oriented to collect data about general knowledge of the participants regarding COVID-19 (source of infection, incubation period, the presence of vaccine, effect of temperature on virus, and the most susceptible people to be affected by the disease). The third section was designed to evaluate the knowledge about COVID-19 symptoms. The fourth section was about the route of disease transmission and the precautions that should be used to prevent or decrease the virus transmission after sneezing. Fifth section was to evaluate the public awareness about the preventive measurements that should be followed by the people. Sixth section was related to the knowledge of the participants about the general treatment of COVID-19. The last section was about the education methods that the participants depend on during the last months of COVID-19 pandemic.

The average time needed to complete the questionnaire was 4 minutes. The questionnaire components were designed based on the educational instructions published by the World health organization (WHO)²².

Assessment of total knowledge: for the 2nd, 3rd, 4th, 5th, and 6th sections, the respondent was given a score =1 for correct answer and zero score for incorrect answer to end up with a total score ranging from 0-35.

Results

Subjects' demographic data (Table 1):

A total of 360 (223 females) subjects participated; most of them were in the age group of 17 to 30 years (65%). 49% were undergraduates and 56% were living in cities. Table 1 shows the percentages of subjects' demographic data.

General knowledge about COVID-19 and the practices to overcome it (Table 2):

The majority of the participants (95%) answered that they have knowledge about COVID-19. Of all participants, 93% know that the incubation period for COVID-19 is about two weeks; 61% believed that the high temperature in summer might kill the virus and 24% answered that the high temperature in summer do not kill the virus. Of all participants, 65% believed that there is no vaccine for COVID-19 but 28% answered that there is a vaccine for COVID-19; 67% believed that bats are a source for COVID-19 and 21% answered that bats are not a source for COVID-19. Regarding the most susceptible and affected groups to infection with the corona virus, the majority of the participants (317) responded that the most susceptible groups are people of all ages who suffer from chronic medical diseases, while (344) indicated that elderly are the most susceptible to infection, and only (97) considered that children are among the most susceptible to infection with SARS-COV-2, and finally (275) indicated that pregnant women are highly susceptible to the infection.

Knowledge about the disease symptoms (Table 3):

Most of participants had good level of knowledge about the symptoms of COVID-19. Most of them (345) know that difficulty of breathing is a symptom for COVID-19, the other symptoms known by them were fever (328), sore throat (326), loss of smell and taste (280), diarrhea (227), runny nose (87), and vomiting (67).

Knowledge about methods of transmission of the disease (Table 4):

Most of the participants (358) know that the disease is transmitted through droplets produced when infected persons cough or sneeze, while (345) believed that the infection with COVID-19 is transmitted through direct contact with contaminated surfaces and then touching eye or mouth. In addition, (149) believed that the infection could be transmitted by eating uncooked meat, (96) believed that the infection transmitted from cats and camels to humans. (63) considered that the infection transmits by insects such as mosquitoes.

Regarding the way by which the participants manage cough or sneezing, 303 participants answered that they use the elbow attachment by arm, (348) use tissue paper to sneeze and cough and get rid of it quickly, (325) answered that they wash hands with warm water and soap after coughing or sneezing, (91) use the hand to cover the mouth, and (8) answered that they don't cover their mouth.

Table 1- Subjects demographic data

characteristic	percentage	No.
Gender		
Male	38%	137
female	62%	223
Age in years		
under 17 years	1%	2
17 -30 years	65%	235
30 – 40 years	31%	110
>40 years	4 %	13
educational level		
Undergraduate	49%	178
Postgraduate	42%	152
Master/Ph. D.	8%	30
marital status		
Unmarried	58%	208
Married	43%	143
current place of residence		
city	56%	159
country	44%	201

Knowledge about the prevention considerations of COVID-19 (Table 5):

Most of participants had good knowledge about prevention considerations that can reduce the disease spread. Most of the participants (344) know that people suffering from Cold, flu, or cough symptoms could be infected with the corona virus and they avoid direct contact with them, (332) answered that they avoid contact with eyes, nose, and mouth before washing their hands and (184) answered that they avoid exposure to animals during travel. 94% believed that children and young adults must take all considerations to prevent infection with COVID-19 although, they are less suspected to be infected. In contrast, 29 of participants believed that medical mask isn't required during travel, (81) revealed that, it is not necessary to wash hands while traveling. Regarding the major steps taken by participants to prevent spread of the disease, most of them (344) believed that personal hygiene and washing hands with water and soap are of the major steps to prevent COVID-19, (327) agreed that following the etiquette of coughing and sneezing is one of the steps also, (304) considered that wearing a medical mask in public places is also one of the steps, (309) agreed to avoid close contact with someone who has cold or flu symptoms, and (185) answered that avoiding unsafe handling of animals is of the steps to prevent COVID-19. In addition, 92% of them agreed that quarantining people infected with the COVID-19 is an effective way to prevent the spread of the virus, while 7% did not agree.. In contrast, (22) believed that taking antibiotics is of the steps to prevent COVID-19.

Knowledge about the treatment and practices of the infected patient (Table 6):

The fact that antibiotics have no role in COVID-19 infection treatment was known by large percent of the participants (82%). In contrast, 6% of them believed that antibiotics are effective in preventing the infection and about 4% of them believed that antibiotics are effective in treating the infection. Regarding the action taken in case of presence of COVID-19 like symptoms, 187 participants indicated that they will call 105 (the hotline for COVID-19 in Egypt), (228) answered that they will go to the nearest emergency hospital and (236) revealed that they will stay at home in a complete isolation.

Also. A few percent (23 participants) of them revealed that they will take medications like antibiotic or chloroquine when they feel symptoms of the COVID-19, while (8) answered that they will not take any action or take to any one, while (84 participants) indicated that they will take medicinal herbs to strengthen their immune system.

Table 2- General knowledge about COVID-19 and the practices to overcome it:

question	No.	%
1- Are you have knowledge about coronavirus?		
• yes	342	95%
• no	18	5%
2- What is the incubation period for Corona virus?		
• Two days	5	1%
• One week	20	6%
• Two weeks	335	93%
3- Is there a vaccine for corona virus?		
• yes	101	28%
• no	234	65%
• I don't know	25	7%
4- Are bats a source of corona virus?		
	242	67%
• yes	75	21%
• no	43	12%
• I don't know		
5- People most susceptible to infection with coronavirus		
• People of all ages who suffer from chronic medical diseases	317	
• children	97	
• Elderly	344	
• Pregnant women	275	
	5	
• I don't know		
6- The high temperature in summer may kill the virus.		
• yes	218	61%
• no	87	24%
• I don't know	55	15%

Determining the most effective method for enhancing the public awareness about the disease (Figure 1):

Social media became the main source of gathering data about COVID-19 according to the response of the participants, followed by TV reports. Participants that depend on scientific publication and WHO reports are a few percent compared to all participants and most of them are master and PhD holders. In addition, other sources for gathering knowledge about the current pandemic were from the direct communication especially with healthcare professionals.

Total knowledge:

The mean \pm SD score of this survey was 27.45 ± 1.2 regarding the knowledge about the disease between the educated people. This indicates that the respondents had good knowledge about the disease and the practices to deal with it. The mean score for females was 27.67 and 27.23 for males.

Discussion

The distribution of the demographic data of the participants showed a high percentage of females, colleague students, unmarried and city dwellers. However, there was no significant difference regarding the total score of both genders.

Knowing the causes is considered the first step in public education²³⁻²⁵. It was shown in previous studies that when people understand how the disease is transmitted, what are the disease symptoms and what the preventive methods for spreading the COVID-19 decrease²⁶⁻²⁸.

Most of the respondents have a good knowledge about COVID-19, but about one-quarter of the participants believed that children are most susceptible to be infected with COVID-19, which indicates that they need more awareness about the disease, however, this point could has a positive impact through following the correct instructions for preventing COVID-19 spread by children.²⁹

Table 3- Knowledge of disease symptoms:

question	No.
1- The symptoms of Coronaviruses	326
• Sore throat	328
• Fever	227
• Diarrhea	345
• difficulty breathing	280
• loss of smell and taste	87
• runny nose	67
• vomiting	1
• I don't know	

Table 4- Knowledge of methods of transmission of the disease:

question	No.
1- How people become infected with Corona virus	358
• Through droplets produced during coughing or sneezing of infected persons.	345
• Through direct contact with contaminated surfaces and then touching eye or mouth.	63
• By insects such as mosquitoes	96
• From cats and camels to humans	149
• By eating uncooked meat	2
• I don't know	
2- Managing cough and sneezing to reduce infection	303
• Use the elbow attachment by arm	348
• Use tissue paper to sneeze and cough and get rid of it quickly	325
• Hand washing with warm water and soap	91
• Use the hand to cover the mouth	8
• I don't cover my mouth	

More of the respondents had a good knowledge about the disease, its symptoms and its treatment. However, a previous study in Saudi Arabia found that the information level about COVID-19 among the public was low and the information level was the significant predictor of both the level of concern and precaution²⁶. In another research in Al Qassim Area³⁰, 44% of the respondents had low knowledge about the source of COVID-19. That might be because the epidemic although in Saudi Arabia in 2013 was limited to a certain area not pandemic like the one we are discussing here. Most of the participant here have a good knowledge about the transmission methods of COVID-19. More than 100 participant depended on the direct communication especially with health care professionals for gathering information about the COVID-19, hence educated people and the healthcare workers should be informed and educated about the new researches and information about COVID-19 to transfer this information to the public in order to reduce the spread of infection spread. Highly educated people (Master and PhD) depend on scientific publications to enhance their knowledge about the disease, however they represent a low percent among the participant.

A lot of the participants answered that when they feel with COVID-19 symptoms, they will call 105 (the hotline for COVID-19 in Egypt). While most of the respondents answered that, they will stay at home with complete isolation or go to the nearest emergency. That indicates the great awareness between the educated people about the correct action that should be followed.

A lot of the participants indicated that, they will not take any antibiotics or take medicines like chloroquine without asking a doctor, as 82% of the participants known that antibiotics has no role in COVID-19 infection, which indicates the high awareness of the participants; the same result in the treatment part was reported in the electronically conducted national survey³¹. The low percent that decided to use antibiotic as a prophylactic medicine could lead to antibiotic misuse, hence, more instructions should be provided by the community pharmacist about the indication for using antibiotic for managing COVID-19.

Table 5- Knowledge of preventive considerations of COVID-19:

question	%	No.
1- Recommendations for travelers during travel		344
<ul style="list-style-type: none"> Avoid approaching people who have cold, flu or cough symptoms No medical mask is required It is not necessary to wash hands while traveling Avoid contact with eyes, nose, and mouth before washing hands Avoid exposure to animals 		29 81 332 184
2- It is not necessary for children and young adults to take any measures to prevent infection with the COVID-19 because they are less likely to be infected		
<ul style="list-style-type: none"> yes no I don't know 	5% 94% 1%	17 340 3
3- Major steps to prevent COVID-19.		
<ul style="list-style-type: none"> Personal hygiene and washing hands with water and soap Following the etiquette of coughing and sneezing Taking antibiotics Wearing a medical mask in public places Avoid close contact with someone who has cold or flu symptoms Avoid unsafe handling of animals 		344 327 22 304 309 185
4- Quarantining people infected with the COVID-19 is an effective way to prevent the spread of the virus		
<ul style="list-style-type: none"> yes no I don't know 	92% 7% 1%	330 25 5

Table 6- Knowledge about treatment and practices of the infected patient:

question	%	NO.
1-Antibiotics		
<ul style="list-style-type: none"> Effective in preventing infection with corona virus Effective for treating corona virus infection It has no role in corona virus infection I don't know 	6% 4% 82% 8%	21 15 307 31
2- What will you do when you feel symptoms of the Corona virus?		
<ul style="list-style-type: none"> Not telling anyone taking any antibiotics or take medicines like chloroquine Buy and eat medicinal herbs to strengthen the immune system you will go to the nearest emergency you will stay at home with complete isolation you will call 105 (the hotline for Coronavirus in Egypt) 		8 23 84 228 236 187




Figure 1- effective method for enhancing the public awareness about the disease

Future information and education messages for enhancing the awareness of the highly educated people should be announced through the main sources for obtaining these data such as social media site, as indicated by the findings of the study most of educated people depend on the social media sites for gathering information about the disease. Also, enhancing the knowledge of educated peoples could result in enhancing the public awareness of the disease through their direct communications with the public. We reported that higher educated people have the ability to gain knowledge about the disease from different sources, which cannot be easily gained by low educated ones.³¹

There was near unanimous approval between the participants on the importance for young adults and children to take all measures to prevent infection with COVID-19 although; they are less likely to be infected. They also agreed on quarantining people infected with the COVID-19 is an effective way to prevent the spread of the virus. That indicates the highly carefulness of the participants from the disease.

Conclusion:

The educated participant had a good knowledge about COVID-19. With recent spreading of this pandemic disease in the world, it is important for the Ministry of Information to enhance the knowledge and awareness of COVID-19. Continuous training is needed by the Ministry of Health to the healthcare workers to support their important role in spreading awareness about COVID-19 infection to the public. Providing the awareness data through the social media could enhance the data delivery to young educated youth. More efforts are needed to enhance the knowledge of the public about the correct use of medications, particularly antibiotics.

References

1. Al-Tawfiq, J.A., Zumla, A., Gautret, P., et al., *Surveillance for emerging respiratory viruses*. The Lancet Infectious Diseases, 2014. **14**(10): p. 992-1000.
2. Fahmy, A.M., Elgendy, M.O., Khalaf, A.M., et al., *COVID-19 Patients with Hepatic Complications During the Third Wave of Pandemic in Egypt*. Journal of Clinical and Nursing Research, 2022. **6**(3): p. 108-121.
3. Mohamed, M.B. and Mahmoud, M.A., *Effects of Educational Program on MERS-Corona-Virus among Nurses Students at Jazan University 1439-2017*. 2018.
4. Elgendy, M.O., Khalaf, A.M., El-Gendy, A.O., et al., *An Observational Study on the Management of COVID-19 Patients in Limited-Resource Hospitals*. Journal of Clinical and Nursing Research, 2022. **6**(3): p. 43-53.
5. Mackay, I.M. and Arden, K.E., *MERS coronavirus: diagnostics, epidemiology and transmission*. Virology journal, 2015. **12**(1): p. 222.
6. Mackay, I.M. and Arden, K.E., *Middle East respiratory syndrome: an emerging coronavirus infection tracked by the crowd*. Virus research, 2015. **202**: p. 60-88.

7. Abdou, L.M., El-Gendy, A.O., Elgendy, M.O., et al., *The Impact of Combining Cefepime or Ceftazidime with Steroidal and Anticoagulant Therapy in the Treatment of COVID-19 Patients*. *NeuroQuantology*, 2022. **20**: p. 3696-3701.
8. Alghamdi, I.G., Hussain, I.I., Almalki, S.S., et al., *The pattern of Middle East respiratory syndrome coronavirus in Saudi Arabia: a descriptive epidemiological analysis of data from the Saudi Ministry of Health*. *International journal of general medicine*, 2014. **7**: p. 417.
9. Battistoni, I., Francioni, M., Morici, N., et al., *Pre-and in-hospital anticoagulation therapy in coronavirus disease 2019 patients: a propensity-matched analysis of in-hospital outcomes*. *Journal of Cardiovascular Medicine*, 2022. **23**(4): p. 264-271.
10. Elgendy, M.O., El-Gendy, A.O., Mahmoud, S., et al., *Side Effects and Efficacy of COVID-19 Vaccines among the Egyptian Population*. *Vaccines*, 2022. **10**(1): p. 109.
11. Zawbaa, H.M., Osama, H., El-Gendy, A., et al., *Effect of mutation and vaccination on spread, severity, and mortality of COVID-19 disease*. *Journal of Medical Virology*, 2021.
12. Elgendy, M.O. and Abdelrahim, M.E., *Public awareness about coronavirus vaccine, vaccine acceptance, and hesitancy*. *Journal of Medical Virology*, 2021.
13. Elgendy, M.O., El-Gendy, A.O., Alzarea, A.I., et al., *SARS-CoV-2 Post Vaccinated Adverse Effects and Efficacy in the Egyptian Population*. *Vaccines*, 2022. **10**(1): p. 18.
14. Eid, R.A., Elgendy, M.O., El-Gendy, A.O., et al., *Efficacy of Ceftazidime and Cefepime in the Management of COVID-19 Patients: Single Center Report from Egypt*. *Antibiotics*, 2021. **10**(11): p. 1278.
15. Jin, Y.-H., Cai, L., Cheng, Z.-S., et al., *A rapid advice guideline for the diagnosis and treatment of 2019 novel coronavirus (2019-nCoV) infected pneumonia (standard version)*. *Military Medical Research*, 2020. **7**(1): p. 4.
16. Elgendy, M.O., Abdelrahman, M.A., Osama, H., et al., *Role of repeating quarantine instructions and healthy practices on COVID-19 patients and contacted persons to raise their awareness and adherence to quarantine instructions*. *International Journal of Clinical Practice*, 2021: p. e14694.
17. Cherkaoui, M., *The Shifting Geopolitics of Coronavirus and the Demise of Neoliberalism—(Part 2)*. Aljazeera Center for Studies, march, 2020. **19**.
18. Elgendy, M.O., Abd Elmawla, M.N., Abdel Hamied, A.M., et al., *COVID-19 patients and contacted person awareness about home quarantine instructions*. *International Journal of Clinical Practice*, 2021. **75**(4): p. e13810.
19. Sayed, A.M., Khalaf, A.M., Abdelrahim, M.E., et al., *Repurposing of some anti-infective drugs for COVID-19 treatment: A surveillance study supported by an in silico investigation*. *International Journal of Clinical Practice*, 2021. **75**(4): p. e13877.
20. Elgendy, M.O., El-Gendy, A.O., and Abdelrahim, M.E., *Public awareness in Egypt about COVID-19 spread in the early phase of the pandemic*. *Patient education and counseling*, 2020. **103**(12): p. 2598-2601.
21. Wang, Y., Di, Y., Ye, J., et al., *Study on the public psychological states and its related factors during the outbreak of coronavirus disease 2019 (COVID-19) in some regions of China*. *Psychology, Health & Medicine*, 2020: p. 1-10.
22. Elbur, A., Alharthi, A., Aljuaid, A., et al., *Knowledge of Middle East Respiratory Syndrome Coronavirus (MERS-CoV) and its Management: A Survey among Saudi People in Taif; Kingdom of Saudi Arabia*. *IOSR Journal Of Pharmacy*, 2016. **6**(2): p. 33-39.
23. Rabbani, S.A., Mustafa, F., and Mahtab, A., *Middle East Respiratory Syndrome (MERS): Awareness among Future Health Care Providers of United Arab Emirates*. *International Journal of Medicine and Public Health*, 2020. **10**(1).
24. Elgendy, M.O., El-Gendy, A.O., Elgendy, S.O., et al., *Perceptions, Knowledge, and Experiences of Using Face Masks among Egyptian Healthcare Workers during the COVID-19 Pandemic: A Cross-Sectional Study*. in *Healthcare*. 2023. MDPI.
25. Zaki, A., Elgendy, M.O., Abdelrahman, M.A., et al., *The Efficacy of Using Different Antibiotics to Prevent Maternal Surgical Site Infections in COVID-19-Infected Cases*.

26. Momattin, H., Mohammed, K., Zumla, A., et al., *Therapeutic options for Middle East respiratory syndrome coronavirus (MERS-CoV)—possible lessons from a systematic review of SARS-CoV therapy*. International Journal of Infectious Diseases, 2013. **17**(10): p. e792-e798.
27. Boshra, M.S., Elgendy, M.O., Abdelaty, L.N., et al., *Knowledge, Attitude, and Acceptance of Sinopharm and AstraZeneca's COVID-19 Vaccines among Egyptian Population: A Cross-Sectional Study*. International Journal of Environmental Research and Public Health, 2022. **19**(24): p. 16803.
28. Shaban, M., Elgendy, M.O., Fahmy, A.M., et al., *The Outcomes of COVID-19 Patients with Spontaneous Intracerebral Hemorrhage Comorbidity and the Efficacy of Enoxaparin in Decreasing the Mortality Rate in Them: Single Egyptian Center Report*. Journal of Personalized Medicine, 2022. **12**(11): p. 1822.
29. Elgendy, S.O., Elgendy, M.O., El-Gendy, A.O., et al., *Health Care Workers' Awareness about the Post-COVID Syndrome and Different Types of COVID-19 Vaccines in Egypt*. NeuroQuantology, 2022. **20**(11): p. 3830-3839.
30. Al-Mohrej, O.A., Al-Shirian, S.D., Al-Otaibi, S.K., et al., *Is the Saudi public aware of Middle East respiratory syndrome?* Journal of infection and public health, 2016. **9**(3): p. 259-266.