

A meta-analysis of the impact on gastrectomy versus endoscopic submucosal dissection for early stomach cancer

¹Rajesh K. Singh

*Correspondence:
Rajesh K. Singh
Department of Pharmaceutical
Chemistry, Shivalik College of
Pharmacy, Nangal, Ropar, 140126,
Punjab, India
Email: rk Singh244@gmail.com

¹ Department of Pharmaceutical
Chemistry, Shivalik College of
Pharmacy, Nangal, Ropar, 140126,
Punjab, India

Volume number 1
Issue number 3
Pages 88-99

Abstract

Background:

We conducted a meta-analysis to assess the impact of gastrectomy versus endoscopic submucosal dissection for early stomach cancer.

There are now a variety of viewpoints on gastrectomy versus endoscopic submucosal dissection for early stomach cancer, and there are not many thorough assessments that are pertinent.

Methods:

A systematic literature search up to July-2022 was performed and 2456 related studies were evaluated. The chosen studies comprised 15461 early stomach cancer subjects who participated in the selected studies' baseline trials; 6503 of them used the endoscopic submucosal dissection, while 8958 used gastrectomy. Odds ratio (OR), and mean difference (MD) with 95% confidence intervals (CIs) were calculated to assess the effect of the gastrectomy versus endoscopic submucosal dissection for early stomach cancer by the dichotomous, and contentious methods with a random or fixed effect model.

Results:

The use of endoscopic submucosal dissection resulted in significantly lower 5-year overall survivals (OR, 0.59; 95% CI, 0.45-0.77, $p < 0.001$), lower the 5-year overall survival in propensity score-matched patients (OR, 0.49; 95% CI, 0.41-0.59, $p < 0.001$), higher recurrences (OR, 6.99; 95% CI, 5.03-9.70, $p < 0.001$), and higher synchronous lesion (OR, 7.24; 95% CI, 2.78-18.83, $p < 0.001$), and higher metachronous lesion (OR, 10.05; 95% CI, 6.44-15.67, $p < 0.001$) compared to the gastrectomy for early stomach cancer.

However, no significant difference was found between submucosal dissection and gastrectomy for early stomach cancer in recurrence-free survival (OR, 0.74; 95% CI, 0.54-1.00, $p = 0.05$), disease-free survival (OR, 0.43; 95% CI, 0.16-1.16, $p = 0.10$), and disease-specific survival (OR, 1.05; 95% CI, 0.38-2.89, $p = 0.92$).

Conclusions:

The use of endoscopic submucosal dissection resulted in significantly lower 5-year overall survival, lower 5-year overall survival in propensity score-matched patients, higher recurrences, higher synchronous lesion, and higher metachronous lesion, however, no significant difference was found in recurrence-free survival, disease-free survival, and disease-specific survival compared to the gastrectomy for early stomach cancer. The small number of studies in several comparisons calls for care when analyzing the results.

Keywords: synchronous lesion; endoscopic submucosal dissection; 5-year overall survival; disease-free survival; disease-specific survival; recurrence; recurrence-free survival; and metachronous lesion

Introduction

The third most common cancer fatality (10%) and the fifth most prevalent malignancy are stomach cancers. ^{1, 2} Regardless of lymph node involvement, early stomach cancer is described as carcinoma that is restricted to the mucosa or submucosa. Historically, the sole curative method for treating early stomach cancer was a radical

surgical gastrectomy with lymph node dissection.³ Radical surgery, however, has been linked to increased morbidity and mortality as well as a decline in quality of life.^{4,5} Endoscopic mucosal resection and endoscopic submucosal dissection are two components of endoscopic resection. For early stomach cancer that was less than 15mm in size, endoscopic mucosal excision was initially advised.⁶ The accepted criteria for endoscopic mucosal excision currently include less than 2 cm of well-differentiated or moderately differentiated adenocarcinoma that is restricted to the mucosa and shows no signs of ulceration or lymphovascular invasion.⁷ The potential for a limited histological examination, particularly if en-bloc resection was not completed, would be the greatest barrier to the widespread adoption of endoscopic mucosal resection in early stomach cancer. To get over the constraints of endoscopic mucosal resection, endoscopic submucosal dissection was developed. By dissecting the submucosal layer with a needle knife during an endoscopic submucosal dissection, a bigger en-bloc resection can be accomplished. When compared to endoscopic mucosal resection, endoscopic submucosal dissection raised the rates of en bloc, histologically complete, and curative resection while decreasing recurrence.⁸ The proposed enlarged indication for endoscopic submucosal dissection was put forth by Gotoda et al. in 2001.⁹ Four distinct criteria make up the Japanese Stomach cancer Association's enlarged indication: (a) differentiated intramucosal cancer, without ulcerative findings, larger than 2 cm; (b) differentiated intramucosal cancer, with ulcerative findings, larger than 3 cm; (c) undifferentiated intramucosal cancer, without ulcerative findings, smaller than 2 cm; and (d) minimal (500 m from the muscularis mucosa) submucosal invasive cancer, differentiated type, larger than 3 cm.¹⁰⁻¹⁴ Though numerous studies have compared the effects of endoscopic submucosal dissection with those of surgical treatment for early stomach cancer, their findings have been inconsistent and their patient populations for both procedures have been heterogeneous.¹⁰⁻¹⁴ To compare the 5-year overall survival rate, disease-specific survival rate, disease-free survival rate, and recurrence-free survival rate of endoscopic submucosal dissection compared to gastrectomy in the treatment of early stomach cancer, a systematic review and meta-analysis were carried out. The study's objective was to determine how endoscopic submucosal dissection versus gastrectomy might affect early stomach cancer.

Method

Eligibility criteria

To create a summary, the study of the endoscopic submucosal dissection in comparison to gastrectomy was selected. The analysis of the impact of gastrectomy versus endoscopic submucosal dissection for early stomach cancer was the major goal of the study.¹⁵

Information sources

The main goals of the current meta-analysis were to evaluate the influence of various outcomes of gastrectomy versus endoscopic submucosal dissection for early stomach cancer. Every selected study involved humans and in any language. Inclusion was unaffected by study size. The publications list was purged of review articles, comments, and research that didn't offer a way to quantify a connotation. The complete course of the study is shown in Figure 1. The following publications were encompassed in the meta-analysis when the inclusion criteria were encountered:

1. The study was either a controlled trial, observational, prospective, or retrospective study.
2. Early stomach cancer topics made up the intended subjects.
3. The intervention program included gastrectomy with an endoscopic submucosal dissection.
4. The study contrasted the gastrectomy versus endoscopic submucosal dissection for early stomach cancer.

The significance of comparison outcomes was not highlighted in studies, and studies that did not examine the effects of endoscopic submucosal dissection in early stomach cancer subjects, research on early stomach cancer without endoscopic submucosal dissection or gastrectomy, and research on early stomach cancer without endoscopic submucosal dissection were excluded from consideration.

Search strategy

A protocol of search approaches was developed following the PICOS concept, and we characterized it as follows: topics for early stomach cancer, P; Endoscopic submucosal dissection technique is the "intervention" or "exposure," whereas the "comparison" was endoscopic submucosal dissection compared to gastrectomy; 5-year overall survival, 5-year overall survival in propensity score-matched patients, synchronous lesion, metachronous lesion, recurrence, recurrence-free survival, disease-free survival, and disease-specific survival were the "outcomes" and finally there are no restrictions on the study's design.¹⁶

We lead a thorough search of the OVID, Embase, Cochrane Library, PubMed, and Google Scholar databases up until June 2022 using an arrangement of keywords and correlated terms for synchronous lesion, endoscopic submucosal dissection, 5-year overall survival, disease-free survival, disease-specific survival, recurrence, recurrence-free survival, and metachronous lesion as shown in Table 1. To avoid studies that did not show a relationship between the endoscopic submucosal dissection and gastrectomy in early stomach cancer individuals, all the papers that had been used were joined into an EndNote file, replicas were eliminated, and the title and abstracts were reviewed and amended.

Selection process

A technique was developed following the epidemiological declaration, which was thereafter arranged and examined in the form of a meta-analysis.




Figure 1 shows a flowchart of the study process.

Data collection process

The criteria used to gather the data included the last name of the primary author, the study period, the publication year, the nation or region, the populace type, the clinical and management physiognomies, the categories, the qualitative and quantitative assessment technique, the information source, the result assessment, and statistical analysis.¹⁷

Data items

When there were disparate findings from a single study founded on the appraisal of the influence of endoscopic submucosal dissection and gastrectomy in early stomach cancer, we independently collected the data.

Study risk of bias assessment

The author individually evaluated the methodology of the designated articles to ascertain the possibility of bias in each study. The methodological quality was evaluated using the "risk of bias instrument" from the Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0. Each study was sorted according to the appraisal criteria and given one of the three risks of bias itemized below: low: A study was rated as having a low risk of bias if all the quality standards were met; if one or more requirements weren't met or weren't encompassed, a study was rated as having a moderate risk of bias. The study was measured to have a high risk of bias in the case that one or more quality criteria were not met at all or were only partially met. The original article was revised to remove any inconsistencies.

Effect measures

Only studies that reported and assessed the influence of endoscopic submucosal dissection in comparison to

gastrectomy were subjected to sensitivity studies. Sensitivity and subclass analysis was utilized to compare the gastrectomy versus endoscopic submucosal dissection for early stomach cancer.

Table 1. Search Strategy for Each Database

Database	Search strategy
Pubmed	#1 "endoscopic submucosal dissection"[MeSH Terms] OR "early stomach cancer"[All Fields] OR "gastrectomy"[All Fields] OR "recurrence"[All Fields] #2 "recurrence-free survival"[MeSH Terms] OR "disease-free survival"[All Fields] OR "5-year overall survival new"[All Fields] OR "disease-specific survival"[All Fields] #3 #1 AND #2
Embase	'endoscopic submucosal dissection'/exp OR 'early stomach cancer'/exp OR 'gastrectomy'/exp OR 'recurrence' #2 'recurrence-free survival'/exp OR '5-year overall survival'/exp OR 'disease-free survival' #3 #1 AND #2
Cochrane library	(endoscopic submucosal dissection):ti,ab,kw (early stomach cancer):ti,ab,kw OR (gastrectomy) :ti,ab,kw (Word variations have been searched) #2 (recurrence):ti,ab,kw OR (recurrence-free survival):ti,ab,kw OR (5-year overall survival) :ti,ab,kw OR (disease-free survival) :ti,ab,kw (Word variations have been searched) #3 #1 AND #2

Synthesis methods

The current meta-analysis used a random- or fixed-effect model with dichotomous and disputed techniques to compute the odds ratio (OR) and mean difference (MD) with a 95 percent confidence interval (CI). It was decided to calculate the I² index, with a range of 0 to 100%. The values around 0%, 25%, 50%, and 75%, respectively, showed no, low, moderate, and high heterogeneity.¹⁸ However, additional characteristics that show a high degree of similarity between the included studies were also analyzed to confirm the employment of the correct model. The random effect was considered if I² was 50% or above; if I² was less than 50%, the likelihood of employing fixed influence rose.¹⁸ However, additional characteristics that show a high degree of similarity between the included studies were also analyzed to confirm the employment of the correct model. A subclass analysis was completed by stratifying the first evaluation based on the previously specified outcome categories. A p-value of 0.05 was used in the analysis to indicate statistical significance for differences across subcategories.

Reporting bias assessment

Publication bias was assessed both qualitatively and statistically using the Egger regression test and funnel plots that display the logarithm of ORs vs their standard errors (publication bias was considered present if p 0.05).¹⁹

Certainty assessment

Two-tailed tests were used to analyze all p-values. The graphs and statistical analysis were created using Reviewer Manager version 5.3 (The Nordic Cochrane Centre, The Cochrane Collaboration, Copenhagen, Denmark).

Results

From a total of 2456 related research that was examined, 24 articles published between 2012 and 2022 that fit the inclusion criteria and were encompassed in the meta-analysis were selected.²⁰⁻⁴³ Table 2 presents the findings from these studies. 15461 early stomach cancer subjects participated in the selected studies' baseline trials; 6503 of them used the endoscopic submucosal dissection, while 8958 used gastrectomy. There were 40 to 3363 subjects present when the trial first began. 21 studies presented data organized by the 5-year overall survivals, 8 studies presented data organized by 5-year overall survival in propensity score-matched patients, 12 studies presented data organized by recurrences, 8 studies presented data organized by synchronous lesion, 14 studies that

Table 2. Characteristics of the selected studies for the meta-analysis

Study	Country	Total	Endoscopic submucosal dissection	Gastrectomy
Chiu, 2012 ²⁰	China	114	74	40
Park, 2014 ²¹	Korea	225	108	117
Kim, 2014 ²²	Korea	158	107	51
Choi, 2015 ²³	Korea	375	261	114
Ryu, 2016 ²⁴	Korea	225	81	144
Cho, 2016 ²⁵	Korea	461	288	173
Pyo, 2016 ²⁶	Korea	2563	1290	1273
Fukunaga, 2017 ²⁷	Japan	308	181	127
Chang, 2017 ²⁸	Korea	153	74	79
Shin, 2017 ²⁹	Korea	275	175	100
Gong, 2017 ³⁰	Korea	79	40	39
Park, 2018 ³¹	Korea	493	111	382
Jeon, 2018 ³²	Korea	617	342	275
Lee, 2018 ³³	Korea	1823	907	916
Hahn, 2018 ³⁴	Korea	1988	786	1202
Bausys, 2019 ³⁵	Lithuania	260	42	218
Hong, 2020 ³⁶	Taiwan	127	26	101
Guo, 2020 ³⁷	China	40	20	20
Pourmousavi, 2020 ³⁸	USA	3363	786	2577
Ahn, 2021 ³⁹	Korea	711	328	383
Quero, 2021 ⁴⁰	Italy	84	42	42
Hirasawa, 2021 ⁴¹	Japan	144	63	81
Lee, 2022 ⁴²	Korean	238	119	119
Kim, 2022 ⁴³	Korean	637	252	385
	Total	15461	6503	8958

presented data organized by metachronous lesion, 10 studies that presented data organized by recurrence-free survival, 9 studies that presented data organized by disease-free survival and 7 studies that presented data organized by the disease-specific survival.

The use of endoscopic submucosal dissection resulted in significantly lower 5-year overall survivals (OR, 0.59; 95% CI, 0.45-0.77, $p < 0.001$) with high heterogeneity ($I^2 = 75\%$), lower the 5-year overall survival in propensity score-matched patients (OR, 0.49; 95% CI, 0.41-0.59, $p < 0.001$) with moderate heterogeneity ($I^2 = 50\%$), higher recurrences (OR, 6.99; 95% CI, 5.03-9.70, $p < 0.001$) with low heterogeneity ($I^2 = 38\%$), and higher synchronous lesion (OR, 7.24; 95% CI, 2.78-18.83, $p < 0.001$) with no heterogeneity ($I^2 = 0\%$), and higher metachronous lesion (OR, 10.05; 95% CI, 6.44-15.67, $p < 0.001$) with no heterogeneity ($I^2 = 0\%$) compared to the gastrectomy for early stomach cancer as shown in Figures 2-6.

However, no significant difference was found between submucosal dissection and gastrectomy for early stomach cancer in recurrence-free survival (OR, 0.74; 95% CI, 0.54-1.00, $p = 0.05$) with low heterogeneity ($I^2 = 33\%$), disease-free survival (OR, 0.43; 95% CI, 0.16-1.16, $p = 0.10$) with high heterogeneity ($I^2 = 86\%$), and disease-specific survival (OR, 1.05; 95% CI, 0.38-2.89, $p = 0.92$) with high heterogeneity ($I^2 = 94\%$) as shown in Figures 7-9. Stratified models could not be utilized to examine the influence of some factors on comparison outcomes, such as gender, age, and ethnicity, due to the dearth of data on these variables. No indication of publication bias was found ($p = 0.87$) after visual analysis of the funnel plot and quantitative assessments using the Egger regression test. The bulk of the included randomized controlled trials, however, were found to have subpar methodological quality, no bias in selective reporting, and scant outcome data.




Figure 2. The effect's forest plot of endoscopic submucosal dissection vs gastrectomy on the 5-year overall survival outcomes in early stomach cancer subjects




Figure 3. The effect's forest plot of endoscopic submucosal dissection vs gastrectomy on the 5-year overall survival in propensity score-matched patients outcomes in early stomach cancer subjects




Figure 4. The effect's forest plot of endoscopic submucosal dissection vs gastrectomy on the recurrence outcomes in early stomach cancer subjects




Figure 5. The effect's forest plot of endoscopic submucosal dissection vs gastrectomy on the synchronous lesion outcomes in early stomach cancer subjects




Figure 6. The effect's forest plot of endoscopic submucosal dissection vs gastrectomy on the metachronous lesion outcomes in early stomach cancer subjects




Figure 7. The effect's forest plot of endoscopic submucosal dissection vs gastrectomy on the recurrence-free survival outcomes in early stomach cancer subjects




Figure 8. The effect's forest plot of endoscopic submucosal dissection vs gastrectomy on the disease-free survival outcomes in early stomach cancer subjects




Figure 9. The effect's forest plot of endoscopic submucosal dissection vs gastrectomy on the disease-specific survival outcomes in early stomach cancer subjects

Discussion

In the trials used for this meta-analysis, 15461 early stomach cancer subjects participated in the selected studies' baseline trials; 6503 of them used the endoscopic submucosal dissection, while 8958 used gastrectomy.²⁰⁻⁴³ The use of endoscopic submucosal dissection resulted in significantly lower 5-year overall survival, lower 5-year overall survival in propensity score-matched patients, higher recurrences, higher synchronous lesion, and higher metachronous lesion compared to the gastrectomy for early stomach cancer. However, no significant difference was found between submucosal dissection and gastrectomy for early stomach cancer in recurrence-free survival, disease-free survival, and disease-specific survival. The small number of studies in several comparisons calls for care when analyzing the results e.g. recurrence-free survival.

Following endoscopic submucosal dissection, there is a greater prevalence of recurring, synchronous, and metachronous lesions. Most typically, the middle or lower portion of the stomach is where primary early gastric malignancies first appear. With a distal gastrectomy, the entire high-risk section of the stomach is removed, leaving just the lower-risk portion. Additionally, endoscopic submucosal dissection enables the persistence of intestinal metaplasia and atrophic gastritis in the remaining mucosa after the procedure.⁴⁴ The 5-year overall survival rate would not be negatively impacted by repeating the endoscopic submucosal dissection if metachronous early gastric tumors are discovered after the procedure. In addition, endoscopic submucosal dissection is less intrusive than surgery, leading to a superior quality of life. Therefore, if metachronous lesions are found early and removed, the increased quality of life with endoscopic submucosal dissection over surgery may offset the minor risk of those lesions. Our findings support the earlier meta-analysis's finding that endoscopic resection had a greater recurrence and metachronous cancer rates than gastrectomy.^{45, 46} In terms of en-bloc resection and recurrence rates, multiple pieces of evidence show that endoscopic submucosal dissection is preferable to endoscopic mucosal resection.^{47, 48} Bleeding and perforation are the primary endoscopic submucosal dissection consequences, and both can be effectively treated intra-procedural in skilled hands.⁴⁹ However, anastomotic leakage, intestinal obstruction, and anastomotic stricture are among the surgical complications that frequently have a considerably greater influence on patients' quality of life, length of hospital stay, and mortality.^{50, 51} Few studies met the inclusion requirements. Some studies omitted descriptions of the random allocation technique, allocation concealment, or blinding. Due to the significant likelihood of bias and the generally poor quality of the papers, the results were not very strong. The study's general conclusions were unaffected by a sensitivity analysis. To collect pertinent research data more thoroughly, improve the standard of the study, and provide reliable and accurate results, randomized controlled trials should be done precisely following methodological principles going ahead. Furthermore, there is a limited

amount of published research on the simultaneous use of studies on endoscopic submucosal dissection and gastrectomy for early stomach cancer. Smaller control and intervention groups were utilized in the majority of the randomized controlled studies included in this study. We believe that these problems could be solved over time and with more research.

This meta-analysis demonstrated how endoscopic submucosal dissection and gastrectomy for early stomach cancer. More research is still needed to clarify these potential connections and compare the impact of endoscopic submucosal dissection and gastrectomy for early stomach cancer on the outcomes under discussion. Larger, more homogeneous samples are required for this investigation. This was also emphasized in a previous study that employed a related meta-analysis technique and found comparable advantageous outcomes for endoscopic submucosal dissection and gastrectomy for early stomach cancer.⁵²⁻⁵⁷ Because our meta-analysis study was unable to determine whether differences in gender, age, and ethnicity are related to the outcomes, well-conducted randomized controlled trials are required to evaluate these factors as well as the combination of different gender, ages, ethnicities, and other variants of subjects.

In conclusion, the use of endoscopic submucosal dissection resulted in significantly lower 5-year overall survival, lower 5-year overall survival in propensity score-matched patients, higher recurrences, higher synchronous lesion, and higher metachronous lesion compared to the gastrectomy for early stomach cancer. However, no significant difference was found between submucosal dissection and gastrectomy for early stomach cancer in recurrence-free survival, disease-free survival, and disease-specific survival.

Limitations

Because several of the studies included in this study were not encompassed in the meta-analysis, there may have been selection bias. The removed publications, nevertheless, did not encounter the requirements for inclusion in our meta-analysis. Furthermore, we were unable to determine whether factors such as age, gender, or ethnicity affected the outcomes. The study aims to compare the outcomes of the gastrectomy group and the endoscopic submucosal dissection group for early stomach cancer. The incorporation of data from earlier studies could have added bias due to incomplete or inaccurate data. Potential sources of bias included the nutritional status of the participants as well as their age and gender characteristics. Unfortunately, certain unpublished papers and missing data can bias the effect being studied.

Conclusions

The use of endoscopic submucosal dissection resulted in significantly lower 5-year overall survival, lower 5-year overall survival in propensity score-matched patients, higher recurrences, higher synchronous lesion, and higher metachronous lesion compared to the gastrectomy for early stomach cancer. However, no significant difference was found between submucosal dissection and gastrectomy for early stomach cancer in recurrence-free survival, disease-free survival, and disease-specific survival. The small number of studies in several comparisons calls for care when analyzing the results e.g. recurrence-free survival.

References

1. Ferlay, J., E. Steliarova-Foucher, J. Lortet-Tieulent, et al. *Cancer incidence and mortality patterns in Europe: estimates for 40 countries in 2012*. European journal of cancer **49**(6): p. 1374-1403 (2013).
2. Ferlay, J., M. Colombet, I. Soerjomataram, et al. *Cancer incidence and mortality patterns in Europe: Estimates for 40 countries and 25 major cancers in 2018*. European journal of cancer **103**: p. 356-387 (2018).
3. Hiki, Y. *Endoscopic mucosal resection (EMR) for early gastric cancer*. Nihon Geka Gakkai Zasshi **97**(4): p. 273-278 (1996).
4. Bonenkamp, J., J. Hermans, M. Sasako, et al. *Extended lymph-node dissection for gastric cancer*. New England Journal of Medicine **340**(12): p. 908-914 (1999).
5. Hartgrink, H., C. Van de Velde, H. Putter, et al. *Extended lymph node dissection for gastric cancer: who may benefit? Final results of the randomized Dutch gastric cancer group trial*. (2004).
6. Ono, H., H. Kondo, T. Gotoda, et al. *Endoscopic mucosal resection for treatment of early gastric cancer*. Gut **48**(2): p. 225-229 (2001).
7. Tsujitani, S., S. Oka, H. Saito, et al. *Less invasive surgery for early gastric cancer based on the low probability of lymph node metastasis*. Surgery **125**(2): p. 148-154 (1999).
8. Oka, S., S. Tanaka, I. Kaneko, et al. *Advantage of endoscopic submucosal dissection compared with EMR for early gastric cancer*. Gastrointestinal endoscopy **64**(6): p. 877-883 (2006).
9. Gotoda, T., A. Yanagisawa, M. Sasako, et al. *Incidence of lymph node metastasis from early gastric cancer: estimation with a large number of cases at two large centers*. Gastric cancer **3**(4): p. 219-225 (2000).
10. Degiuli, M. and F. Calvo *Survival of early gastric cancer in a specialized European center. Which lymphadenectomy is necessary?* World journal of surgery **30**(12): p. 2193-2203 (2006).
11. Ishikawa, S., A. Togashi, M. Inoue, et al. *Indications for EMR/ESD in cases of early gastric cancer: relationship between histological type, depth of wall invasion, and lymph node metastasis*. Gastric Cancer **10**(1): p. 35-38 (2007).

12. Takenaka, R., Y. Kawahara, H. Okada, et al. *Risk factors associated with local recurrence of early gastric cancers after endoscopic submucosal dissection*. *Gastrointestinal endoscopy* **68**(5): p. 887-894 (2008).
13. Lian, J., S. Chen, Y. Zhang, et al. *A meta-analysis of endoscopic submucosal dissection and EMR for early gastric cancer*. *Gastrointestinal endoscopy* **76**(4): p. 763-770 (2012).
14. Uedo, N., H. Iishi, M. Tatsuta, et al. *Longterm outcomes after endoscopic mucosal resection for early gastric cancer*. *Gastric Cancer* **9**(2): p. 88-92 (2006).
15. Stroup, D.F., J.A. Berlin, S.C. Morton, et al. *Meta-analysis of observational studies in epidemiology: a proposal for reporting*. *Jama* **283**(15): p. 2008-2012 (2000).
16. Liberati, A., D.G. Altman, J. Tetzlaff, et al. *The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration*. *Journal of clinical epidemiology* **62**(10): p. e1-e34 (2009).
17. Gupta, S., G. Rout, A.H. Patel, et al. *Efficacy of generic oral directly acting agents in patients with hepatitis C virus infection*. *Journal of viral hepatitis* **25**(7): p. 771-778 (2018).
18. Sheikhabahaei, S., T.J. Trahan, J. Xiao, et al. *FDG-PET/CT and MRI for evaluation of pathologic response to neoadjuvant chemotherapy in patients with breast cancer: a meta-analysis of diagnostic accuracy studies*. *The oncologist* **21**(8): p. 931-939 (2016).
19. Higgins, J.P., S.G. Thompson, J.J. Deeks, et al. *Measuring inconsistency in meta-analyses*. *Bmj* **327**(7414): p. 557-560 (2003).
20. Chiu, P.W.Y., A.Y.B. Teoh, K.F. To, et al. *Endoscopic submucosal dissection (ESD) compared with gastrectomy for treatment of early gastric neoplasia: a retrospective cohort study*. *Surgical endoscopy* **26**(12): p. 3584-3591 (2012).
21. Park, C.H., H. Lee, D.W. Kim, et al. *Clinical safety of endoscopic submucosal dissection compared with surgery in elderly patients with early gastric cancer: a propensity-matched analysis*. *Gastrointestinal endoscopy* **80**(4): p. 599-609 (2014).
22. Kim, D.Y., S.J. Hong, G.S. Cho, et al. *Long-term efficacy of endoscopic submucosal dissection compared with surgery for early gastric cancer: a retrospective cohort study*. *Gut and liver* **8**(5): p. 519 (2014).
23. Choi, I.J., J.H. Lee, Y.-I. Kim, et al. *Long-term outcome comparison of endoscopic resection and surgery in early gastric cancer meeting the absolute indication for endoscopic resection*. *Gastrointestinal endoscopy* **81**(2): p. 333-341. e1 (2015).
24. Ryu, S.J., B.-W. Kim, B.G. Kim, et al. *Endoscopic submucosal dissection versus surgical resection for early gastric cancer: a retrospective multicenter study on immediate and long-term outcome over 5 years*. *Surgical endoscopy* **30**(12): p. 5283-5289 (2016).
25. Cho, J.-H., S.-W. Cha, H.G. Kim, et al. *Long-term outcomes of endoscopic submucosal dissection for early gastric cancer: a comparison study to surgery using propensity score-matched analysis*. *Surgical endoscopy* **30**(9): p. 3762-3773 (2016).
26. Pyo, J.H., H. Lee, B.-H. Min, et al. *Long-term outcome of endoscopic resection vs. surgery for early gastric cancer: a non-inferiority-matched cohort study*. *Official journal of the American College of Gastroenterology* *ACG* **111**(2): p. 240-249 (2016).
27. Fukunaga, S., Y. Nagami, M. Shiba, et al. *Long-term prognosis of expanded-indication differentiated-type early gastric cancer treated with endoscopic submucosal dissection or surgery using propensity score analysis*. *Gastrointestinal endoscopy* **85**(1): p. 143-152 (2017).
28. Chang, J.Y., K.-N. Shim, C.H. Tae, et al. *Comparison of clinical outcomes after endoscopic submucosal dissection and surgery in the treatment of early gastric cancer: a single-institute study*. *Medicine* **96**(30) (2017).
29. Shin, D.W., H.Y. Hwang, and S.W. Jeon *Comparison of endoscopic submucosal dissection and surgery for differentiated type early gastric cancer within the expanded criteria*. *Clinical Endoscopy* **50**(2): p. 170-178 (2017).
30. Gong, E.J., D.H. Kim, J.Y. Ahn, et al. *Comparison of long-term outcomes of endoscopic submucosal dissection and surgery for esophagogastric junction adenocarcinoma*. *Gastric Cancer* **20**(1): p. 84-91 (2017).
31. Park, J.C., Y.K. Lee, S.Y. Kim, et al. *Long-term outcomes of endoscopic submucosal dissection in comparison to surgery in undifferentiated-type intramucosal gastric cancer using propensity score analysis*. *Surgical endoscopy* **32**(4): p. 2046-2057 (2018).
32. Jeon, H.K., G.H. Kim, B.E. Lee, et al. *Long-term outcome of endoscopic submucosal dissection is comparable to that of surgery for early gastric cancer: a propensity-matched analysis*. *Gastric Cancer* **21**(1): p. 133-143 (2018).
33. Lee, S., K.D. Choi, M. Han, et al. *Long-term outcomes of endoscopic submucosal dissection versus surgery in early gastric cancer meeting expanded indication including undifferentiated-type tumors: a criteria-based analysis*. *Gastric Cancer* **21**(3): p. 490-499 (2018).
34. Hahn, K.Y., C.H. Park, Y.K. Lee, et al. *Comparative study between endoscopic submucosal dissection and surgery in patients with early gastric cancer*. *Surgical endoscopy* **32**(1): p. 73-86 (2018).

35. Bausys, R., A. Bausys, J. Stanaitis, et al. *Propensity score-matched comparison of short-term and long-term outcomes between endoscopic submucosal dissection and surgery for treatment of early gastric cancer in a Western setting*. *Surgical Endoscopy* **33**(10): p. 3228-3237 (2019).
36. Hong, T.-C., J.-M. Liou, C.-C. Yeh, et al. *Endoscopic submucosal dissection comparing with surgical resection in patients with early gastric cancer—a single center experience in Taiwan*. *Journal of the Formosan Medical Association* **119**(12): p. 1750-1757 (2020).
37. Guo, A., C. Du, S. Tian, et al. *Long-term outcomes of endoscopic submucosal dissection versus surgery for treating early gastric cancer of undifferentiated-type*. *Medicine* **99**(22): p. e20501 (2020).
38. Pourmousavi, M.K., R. Wang, T. Kerdsirichairat, et al. *Comparable cancer-specific mortality of patients with early gastric cancer treated with endoscopic therapy vs surgical resection*. *Clinical Gastroenterology and Hepatology* **18**(12): p. 2824-2832. e1 (2020).
39. Ahn, J.Y., Y.-I. Kim, W.G. Shin, et al. *Comparison between endoscopic submucosal resection and surgery for the curative resection of undifferentiated-type early gastric cancer within expanded indications: a nationwide multi-center study*. *Gastric Cancer* **24**(3): p. 731-743 (2021).
40. Quero, G., C. Fiorillo, F. Longo, et al. *Propensity score-matched comparison of short-and long-term outcomes between surgery and endoscopic submucosal dissection (ESD) for intestinal type early gastric cancer (EGC) of the middle and lower third of the stomach: a European tertiary referral center experience*. *Surgical Endoscopy* **35**(6): p. 2592-2600 (2021).
41. Hirasawa, T., N. Yamamoto, and T. Sano *Is endoscopic resection appropriate for type 3 gastric neuroendocrine tumors? Retrospective multicenter study*. *Digestive Endoscopy* **33**(3): p. 408-417 (2021).
42. Lee, G.H., E. Lee, B. Park, et al. *Long-term outcomes of endoscopic submucosal dissection and surgery for undifferentiated intramucosal gastric cancer regardless of size*. *World Journal of Gastroenterology* **28**(8): p. 840 (2022).
43. Kim, J.-H., Y.-I. Kim, J.Y. Ahn, et al. *Long-term outcomes of endoscopic resection followed by additional surgery after non-curative resection in undifferentiated-type early gastric cancer: a nationwide multi-center study*. *Surgical Endoscopy* **36**(3): p. 1847-1856 (2022).
44. Choi, Y.K., J.Y. Ahn, H.K. Na, et al. *Outcomes of endoscopic submucosal dissection for gastric epithelial neoplasm in chronic kidney disease patients: propensity score-matched case-control analysis*. *Gastric Cancer* **22**(1): p. 164-171 (2019).
45. Kondo, A., E.G.H. de Moura, W.M. Bernardo, et al. *Endoscopy vs surgery in the treatment of early gastric cancer: systematic review*. *World Journal of Gastroenterology* **21**(46): p. 13177 (2015).
46. Meng, F.-S., Z.-H. Zhang, Y.-M. Wang, et al. *Comparison of endoscopic resection and gastrectomy for the treatment of early gastric cancer: a meta-analysis*. *Surgical endoscopy* **30**(9): p. 3673-3683 (2016).
47. Ning, F.-L., C.-D. Zhang, P. Wang, et al. *Endoscopic resection versus radical gastrectomy for early gastric cancer in Asia: A meta-analysis*. *International Journal of Surgery* **48**: p. 45-52 (2017).
48. Park, Y.-M., E. Cho, H.-Y. Kang, et al. *The effectiveness and safety of endoscopic submucosal dissection compared with endoscopic mucosal resection for early gastric cancer: a systematic review and metaanalysis*. *Surgical endoscopy* **25**(8): p. 2666-2677 (2011).
49. Farhat, S., S. Chaussade, T. Ponchon, et al. *Endoscopic submucosal dissection in a European setting. A multi-institutional report of a technique in development*. *Endoscopy* **43**(08): p. 664-670 (2011).
50. Kitano, S., N. Shiraishi, I. Uyama, et al. *A multicenter study on oncologic outcome of laparoscopic gastrectomy for early cancer in Japan*. *Annals of surgery* **245**(1): p. 68 (2007).
51. Ryu, K.W., Y.-W. Kim, J.H. Lee, et al. *Surgical complications and the risk factors of laparoscopy-assisted distal gastrectomy in early gastric cancer*. *Annals of Surgical Oncology* **15**(6): p. 1625-1631 (2008).
52. Gu, L., P.A. Khadaroo, L. Chen, et al. *Comparison of long-term outcomes of endoscopic submucosal dissection and surgery for early gastric cancer: a systematic review and meta-analysis*. *Journal of Gastrointestinal Surgery* **23**(7): p. 1493-1501 (2019).
53. Benites-Gofñi, H., F. Palacios-Salas, A. Carlin-Ronquillo, et al. *Endoscopic submucosal dissection versus surgery for patients with undifferentiated early gastric cancer*. *Revista Espanola de Enfermedades Digestivas: Organo Oficial de la Sociedad Espanola de Patologia Digestiva* (2022).
54. Abdelfatah, M.M., M. Barakat, D. Ahmad, et al. *Long-term outcomes of endoscopic submucosal dissection versus surgery in early gastric cancer: a systematic review and meta-analysis*. *European journal of gastroenterology & hepatology* **31**(4): p. 418-424 (2019).
55. Liu, Q., L. Ding, X. Qiu, et al. *Updated evaluation of endoscopic submucosal dissection versus surgery for early gastric cancer: A systematic review and meta-analysis*. *International Journal of Surgery* **73**: p. 28-41 (2020).
56. Huh, C.-W., D.W. Ma, B.-W. Kim, et al. *Endoscopic submucosal dissection versus surgery for undifferentiated-type early gastric cancer: a systematic review and meta-analysis*. *Clinical Endoscopy* **54**(2): p. 202-210 (2021).

57. Yang, H.-J., J.-H. Kim, N.W. Kim, et al. *Comparison of long-term outcomes of endoscopic submucosal dissection and surgery for undifferentiated-type early gastric cancer meeting the expanded criteria: a systematic review and meta-analysis*. Surgical endoscopy: p. 1-12 (2022).